

Res. on Crops **14** (3) : 897-901 (2013)

With two figures

Printed in India

A simple and low-cost technique of DNA extraction from edible mushrooms examined by molecular phylogenetics

FARHAT AHMADI AVIN*, SUBHA BHASSU** AND VIKINESWARY SABARATNAM

Mushroom Research Centre (MRC)

University of Malaya, 50603, Kuala Lumpur, Malaysia

**(e-mail : farhat.avin@gmail.com; **subhabhassu@gmail.com)*

(Received : March 2013)

ABSTRACT

The first and most important step of molecular techniques is to isolate the high quality and standard quantity of DNA. The DNA extracted using the recommended method could successfully amplify the regions of interest and demonstrated reliable results can be applied in other molecular assays. Moreover, the designed primers of ITS1-UM2 and ITS4-UM2 were perfectly matched with the species of Basidiomycetes, can be used in phylogenetic studies of other mushrooms. We here evaluated the quality and quantity of DNA using a spectrophotometer, showed reliable OD_{260/280}, and concentration. The protocol is efficient, rapid, low-cost, and simple, needs low amount of sample, and requires minimum facilities. The standard yield in addition to the high quality of DNA will enable mycologists to establish molecular techniques easier. In the current study, the constructed phylogenetic tree based on the obtained sequences of Internal Transcribed Spacer (ITS) I and II regions distinctly classified the examined material.

Key words : Basidiomycetes, DNA isolation, Internal Transcribed Spacer (ITS), phylogenetic, spectrophotometer

ACKNOWLEDGEMENTS

This study was supported by the research grants PS291-2009C, PV088-2011B and 66-02-03-0074 from the Institute of Research Management and Monitoring (IPPP), University of Malaya.

REFERENCES

- Avin, F. A., Bhassu, S., Shin, T. Y. and Sabaratnam, V. (2012) Molecular classification and phylogenetic relationships of selected edible Basidiomycetes species. *Mole. Biol. Reports* **39** : 1-10.
- Bao, D., Kinugasa, S. and Kitamoto, Y. (2004). The biological species of oyster mushrooms (*Pleurotus* spp.) from Asia based on mating compatibility tests. *J. Wood Sci.* **50** : 162-68.
- Choi, D. B., Ding, J. L. and Cha, W. S. (2007). Homology search of genus *Pleurotus* using an internal transcribed spacer region. *Korean J. Chem. Engg.* **24** : 408-12.
- Choi, D. B., Nam, H. G. and Cha, W. S. (2006). Studies on cultivation and biological activities of *Pleurotus nebrodensis* in zeng. *Korean J. Chem. Engg.* **23** : 241-46.
- Eichlerová, I. and Homolka, L. (1999). Preparation and crossing of basidiospore-derived monokaryons—a useful tool for obtaining laccase and other ligninolytic enzyme higher-producing dikaryotic strains of *Pleurotus ostreatus*. *Antonie van Leeuwenhoek* **75** : 321-27.
- Fredricks, D. N., Smith, C. and Meier, A. (2005). Comparison of six DNA extraction methods for recovery of fungal DNA as assessed by quantitative PCR. *J. Clinical Microbiol.* **43** : 5122.
- Gharehaghaji, A., Goltapeh, E., Masiha, S. and Gordan, H. (2007). Hybrid production of oyster mushroom, *Pleurotus ostreatus* (Jacq : Fries) Kummer. *Pak. J. Biol. Sci.* **10** : 2334.
- Ibrahim, R. (2011). A modified CTAB protocol for DNA extraction from young flower petals of some medicinal plant species. *Gene Conserve.* **10** : 165-82.
- Kumar, O. S., Sharma, M. and Singh, D. (2006). DNA isolation from goat blood using different brands of household detergents and its downstream application. *Indian J. Exptl. Biol.* **44** : 852.
- Librado, P. and Rozas, J. (2009). DnaSP v5 : a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* **25** : 1451-52.
- Manian, S., Sreenivasaprasad, S. and Mills, P. (2001). DNA extraction method for PCR in mycorrhizal fungi. *Lett. Appl. Microbiol.* **33** : 307-10.
- P3aza, G., Upchurch, R., Brigmon, R., Whitman, W. and Ulfig, K. (2004). Rapid DNA extraction for screening soil filamentous fungi using PCR amplification. *Polish J. Environ. Studies* **13** : 315-18.
- Ro, H. S., Kim, S. S., Ryu, J. S., Jeon, C. O., Lee, T. S. and Lee, H. S. (2007). Comparative studies on the diversity of the edible mushroom *Pleurotus eryngii*: ITS sequence analysis, RAPD fingerprinting, and physiological characteristics. *Mycol. Res.* **111** : 710-15.
- Rozen, S. and Skaletsky, H. (2000). Primer3 on the WWW for general users and for biologist programmers. *Methods Mol. Biol.* **132** : 365-86.
- Sharma, P., Joshi, N. and Sharma, A. (2010). Isolation of genomic DNA from medicinal plants without liquid nitrogen. *Indian J. Exp. Biol.* **48** : 610-14.
- Shnyreva, A. and Shtaer, O. (2006). Differentiation of closely related oyster fungi *Pleurotus pulmonarius* and *P. ostreatus* by mating and molecular markers. *Russian J. Genet.* **42** : 539-45.
- Sokal, R. R. and Michener, C. D. (1958). A statistical method for evaluating systematic relationships. *Univ. Kans. Sci. Bull.* **38** : 1409-38.
- Stajic, M., Sikorski, J., Wasser, S. P. and Nevo, E. (2005). Genetic similarity and taxonomic relationships within the genus *Pleurotus* (higher Basidiomycetes) determined by RAPD analysis. *Mycotaxon* **93** : 247-56.
- Tamura, K., Dudley, J., Nei, M. and Kumar, S. (2007). MEGA4 : molecular evolutionary genetics analysis (MEGA) software version 4.0. *Mole. Biol. and Evol.* **24** : 1596-99.
- Tsai, Y. L. and Olson, B. H. (1991). Rapid method for direct extraction of DNA from soil and sediments. *Appl. and Environ. Microbiol.* **57** : 1070-74.
- Weiland, J. (1997). Rapid procedure for the extraction of DNA from fungal spores and mycelia. *Fungal Genet. Newsl.* **44** : 60-63.
- White, T., Bruns, T., Lee, S. and Taylor, J. (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *PCR Protocols—A Guide to Methods and Applications.* pp. 315-22.
- Yamada, Y., Makimura, K., Merhendi, H., Ueda, K., Nishiyama, Y., Yamaguchi, H. and Osumi, M. (2002). Comparison of different methods for extraction of mitochondrial DNA from human pathogenic yeasts. *Jap. J. Infect. Dis.* **55** : 122-25.

